

Disclaimer: These solutions are provided by the academic team of BASE. These solutions may be different from those to be provided by the Central Board of Secondary Education (CBSE) due to difference in assumptions taken in support of the solutions. In such cases solution provided by CBSE will be deemed as final. Users are advised to verify/check any information with the relevant Examination Board(s) and/or other source(s), and to obtain any appropriate professional advice before acting on the information provided in the Portal.

Every effort is made to provide accurate and complete information on this website. However, though all efforts have been made to ensure the accuracy, completeness, usefulness or otherwise of the content on this Portal, neither BASE, nor their employees, representatives or Board members make any warranty, expressed or implied, with respect to information available from this website. Additionally, BASE assumes no legal liability for the accuracy, completeness or usefulness of any information disclosed herein.

In no event will the BASE be liable for any expense, loss or damage including, without limitation, indirect or consequential loss or damage, or any expense, loss or damage whatsoever arising from use, or loss of use, of data, arising out of or in connection with the use of this Portal.

NEET-1 (AIPMT) May, 2016 Test Paper Code - B KEYS (Revised)

Q. No.	Chemistry	Q. No.	Botany	Q. No.	Zoology	Q. No.	Physics
1.	(3)	46.	(3)	91.	(4)	136.	(3)
2.	(4)	47.	(3)	92.	(4)	137.	(3)
3.	(4)	48.	(3)	93.	(1)	138.	(3)
4.	(4)	49.	(1)	94.	(2)	139.	(3)
5.	(1)	50.	(2)	95.	(1)	140.	(3)
6.	(3)	51.	(2)	96.	(1)	141.	(3)
7.	(4)	52.	(3)	97.	(2)	142.	(4)
8.	(3)	53.	(2)	98.	(1)	143.	(4)
9.	(4)	54.	(1)	99.	(1)	144.	(3)
10.	(3)	55.	(1)	100.	(4)	145.	(3)
11.	(1)	56.	(4)	101.	(1)	146.	(2)
12.	(4)	57.	(1)	102.	(1)	147.	(1)
13.	(3)	58.	(3)	103.	(4)	148.	(3)

14.	(2)	59.	(1)	104.	–	149.	(1)
15.	(4)	60.	(1)	105.	(2)	150.	(4)
16.	(4)	61.	(2)	106.	(2)	151.	(2)
17.	(1)	62.	(1)	107.	(2)	152.	(3)
18.	(3)	63.	(2)	108.	(4)	153.	(1)
19.	(1)	64.	(3)	109.	(4)	154.	(3)
20.	(3)	65.	(2)	110.	(1)	155.	(4)
21.	(2)	66.	(3)	111.	(3)	156.	(3)
22.	(4)	67.	(3)	112.	(3)	157.	(2)
23.	(1) & (2)	68.	(4)	113.	(1)	158.	(1)
24.	(4)	69.	(1)	114.	(2)	159.	(1)
25.	(4)	70.	(3)	115.	(1)	160.	(1)
26.	(4)	71.	(2)	116.	(2)	161.	(3)
27.	(1)	72.	(1)	117.	(3)	162.	(4)
28.	(4)	73.	(4)	118.	(4)	163.	(1)
29.	(1)	74.	(4)	119.	(4)	164.	(2)
30.	(4)	75.	(3)	120.	(3)	165.	(4)
31.	(3)	76.	(4)	121.	(4)	166.	(2)
32.	(1)	77.	(1)	122.	(2)	167.	(1)
33.	(4)	78.	(2)	123.	(1)	168.	(2)
34.	(1)	79.	(3)	124.	(3)	169.	(2)
35.	(1)	80.	(3)	125.	(3)	170.	(2)
36.	(4)	81.	(3)	126.	(1)	171.	(4)
37.	(4)	82.	(4)	127.	(2)	172.	(3)
38.	(1)	83.	(3)	128.	(2)	173.	(1)
39.	(3)	84.	(1)	129.	(4)	174.	(1)
40.	(4)	85.	(1)	130.	(2)	175.	(4)
41.	(3)	86.	(4)	131.	(4)	176.	(4)
42.	(1)	87.	(3)	132.	(4)	177.	(4)
43.	(2)	88.	(2)	133.	(2)	178.	(2)
44.	(4)	89.	(4)	134.	(1)	179.	(4)
45.	(4)	90.	(2)	135.	(3)	180.	(1)

* * *

NEET-1 (AIPMT) May, 2016


Test Paper Code – B

Questions with Solutions (Revised)

1. The addition of a catalyst during a chemical reaction alters which of the following quantities?

- (1) Internal energy
- (2) Enthalpy
- (3) Activation energy
- (4) Entropy

Ans (3)



2. Predict the correct order among the following:

- (1) lone pair – lone pair > bond pair – bond pair > lone pair – bond pair
- (2) bond pair – bond pair > lone pair – bond pair > lone pair – lone pair
- (3) lone pair – bond pair > bond pair – bond pair > lone pair – lone pair
- (4) lone pair – lone pair > lone pair – bond pair > bond pair – bond pair

Ans (4)

3. The correct statement regarding the basicity of arylamines is

- (1) Arylamines are generally more basic than alkylamines because the nitrogen lone-pair electrons are not delocalized by interaction with the aromatic ring π electron system.
- (2) Arylamines are generally more basic than alkylamines because of aryl group.
- (3) Arylamines are generally more basic than alkylamines, because the nitrogen atom in arylamines is sp -hybridized.
- (4) Arylamines are generally less basic than alkylamines because the nitrogen lone-pair electrons are delocalized by interaction with the aromatic ring π electron system.

Ans (4)




4. When copper is heated with conc. HNO_3 it produces

- (1) $\text{Cu}(\text{NO}_3)_2$ and NO
- (2) $\text{Cu}(\text{NO}_3)_2$, NO and NO_2
- (3) $\text{Cu}(\text{NO}_3)_2$ and N_2O
- (4) $\text{Cu}(\text{NO}_3)_2$ and NO_2

Ans (4)



5. For the following reactions



Which of the following statements is correct?

- (1) (a) is elimination, (b) is substitution and (c) is addition reaction.
- (2) (a) is elimination, (b) and (c) are substitution reactions.
- (3) (a) is substitution, (b) and (c) are addition reactions.
- (4) (a) and (b) are elimination reactions and (c) is addition reaction.

Ans (1)

a = elimination

b = substitution


c = addition

6. Two electrons occupying the same orbital are distinguished by:


- (1) Magnetic quantum number
- (2) Azimuthal quantum number
- (3) Spin quantum number
- (4) Principal quantum number

Ans (3)

According to Pauli's exclusion principle,



7. The reaction



can be classified as

- (1) Alcohol formation reaction
- (2) Dehydration reaction
- (3) Williamson alcohol synthesis reaction
- (4) Williamson ether synthesis reaction


Ans (4)

Williamson ether synthesis for the formation of methoxy cyclopentane.

8. The electronic configuration of Eu(Atomic No.63), Gd(Atomic No.64) and Tb(Atomic No.65) are

- (1) $[\text{Xe}]4f^65d^16s^2$, $[\text{Xe}]4f^75d^16s^2$ and $[\text{Xe}]4f^96s^2$
- (2) $[\text{Xe}]4f^65d^16s^2$, $[\text{Xe}]4f^75d^16s^2$ and $[\text{Xe}]4f^85d^16s^2$
- (3) $[\text{Xe}]4f^76s^2$, $[\text{Xe}]4f^75d^16s^2$ and $[\text{Xe}]4f^96s^2$
- (4) $[\text{Xe}]4f^76s^2$, $[\text{Xe}]4f^86s^2$ and $[\text{Xe}]4f^85d^16s^2$

Ans (3)



9. At 100°C the vapour pressure of a solution of 6.5 g of a solute in 100 g water is 732 mm. If $K_b = 0.52$, the boiling point of this solution will be

- (1) 100°C (2) 102°C (3) 103°C (4) 101°C

Ans (4)

$$W_2 = 6.5 \text{ g}$$

$$W_1 = 100 \text{ g}$$

$$p_1^\circ = 760 \text{ mm of Hg}$$

$$p_1 = 732 \text{ mm}$$

$$K_b = 0.52$$

$$\frac{p_1^\circ - p_1}{p_1^\circ} = \frac{W_2}{M_2} \times \frac{M_1}{W_1}$$

$$M_2 = \frac{p_1^\circ}{p_1^\circ - p_1} \times \frac{W_2 M_1}{W_1}$$

$$M_2 = \frac{760}{(760 - 732)} \times \frac{6.5 \times 18}{100} = 31.75 \text{ g/mol}$$

$$\Delta T_b = \frac{1000 \times k_b \times W_2}{M_2 \times W_1} = \frac{1000 \times 0.52 \times 6.5}{31.75 \times 100}$$

$$\Delta T = 1.06^\circ\text{C}$$

$$\therefore T_b = T_b^\circ + \Delta T_b$$

$$= 100 + 1.06^\circ\text{C} = 101.06^\circ\text{C}$$

$$= 101^\circ\text{C}$$

10. The correct statement regarding the comparison of staggered and eclipsed conformations of ethane, is
- (1) The eclipsed conformation of ethane is more stable than staggered conformation, because eclipsed conformation has no torsional strain.
 - (2) The eclipsed conformation of ethane is more stable than staggered conformation even though the eclipsed conformation has torsional strain
 - (3) The staggered conformation of ethane is more stable than eclipsed conformation, because staggered conformation has no torsional strain
 - (4) The staggered conformation of ethane is less stable than eclipsed conformation, because staggered conformation has torsional strain

Ans (3)

11. Which one of the following characteristics is associated with adsorption?

- (1) ΔG , ΔH and ΔS all are negative
- (2) ΔG and ΔH are negative but ΔS is positive
- (3) ΔG and ΔS are negative but ΔH is positive
- (4) ΔG is negative but ΔH and ΔS are positive

Ans (1)

For adsorption, $\Delta G < 0$ (spontaneous)
 $\Delta H < 0$ (exothermic)
 $\Delta S < 0$ (entropy decreases for the adsorbate)


12. Match the compounds given in column I with the hybridization and shape given in column II and mark the correct option.

Column – I		Column – II	
(a)	XeF ₆	(i)	distorted octahedral
(b)	XeO ₃	(ii)	square planar
(c)	XeOF ₄	(iii)	pyramidal
(d)	XeF ₄	(iv)	square pyramidal

Code


	(a)	(b)	(c)	(d)
(1)	(i)	(ii)	(iv)	(iii)
(2)	(iv)	(iii)	(i)	(ii)
(3)	(iv)	(i)	(ii)	(iii)
(4)	(i)	(iii)	(iv)	(ii)

Ans (4)



13. The correct statement regarding a carbonyl compound with a hydrogen atom on its alpha-carbon, is
- (1) a carbonyl compound with a hydrogen atom on its alpha-carbon rapidly equilibrates with its corresponding enol and this process is known as aldehyde-ketone equilibration.
 - (2) a carbonyl compound with a hydrogen atom on its alpha-carbon rapidly equilibrates with its corresponding enol and this process is known as carbonylation.
 - (3) a carbonyl compound with hydrogen atom on its alpha-carbon rapidly equilibrates with its corresponding enol and this process is known as keto-enol tautomerism.
 - (4) a carbonyl compound with a hydrogen atom on its alpha-carbon never equilibrates with its corresponding enol.

Ans (3)



14. In a protein molecule various amino acids are linked together by
- (1) β -glycosidic bond
 - (2) peptide bond
 - (3) dative bond
 - (4) α -glycosidic bond

Ans (2)

Peptide bonds are found in proteins.

15. Match items with column I with the items of column II and assign the correct code

Column – I		Column – II	
(a)	Cyanide process	(i)	Ultrapure Ge
(b)	Froth Floatation process	(ii)	Dressing of ZnS
(c)	Electrolytic reduction	(iii)	Extraction of Al
(d)	Zone refining	(iv)	Extraction of Au
		(v)	Purification of Ni

Code

	(a)	(b)	(c)	(d)
(1)	(ii)	(iii)	(i)	(v)
(2)	(i)	(ii)	(iii)	(iv)
(3)	(iii)	(iv)	(v)	(i)
(4)	(iv)	(ii)	(iii)	(i)

Ans (4)

16. Which of the following is an analgesic?

(1) Penicillin (2) Streptomycin (3) Chloromycetin (4) Novalgin

Ans (4)


Penicillin, streptomycin and chloromycetin are antibiotics while novalgin is analgesic.

17. Which is the correct statement for the given acids?

- (1) Phosphinic acid is a monoprotic acid while phosphonic acid is a diprotic acid.
 (2) Phosphinic acid is a diprotic acid while phosphonic acid is a monoprotic acid.
 (3) Both are triprotic acids
 (4) Both are diprotic acids


Ans (1)

Phosphinic acid
(hypophosphorous acid)



Basicity = 1

Phosphonic acid
(ortho phosphorous acid)




Basicity = 2

18. The pair of electron in the given carbanion, $\text{CH}_3\text{C}\equiv\text{C}^-$, is present in which of the following orbitals?

(1) sp^3 (2) sp^2 (3) sp (4) $2p$

Ans (3)



19. Consider the molecules CH_4 , NH_3 and H_2O . Which of the given statements is false?

- (1) The H–O–H bond angle in H_2O is larger than the H–C–H bond angle in CH_4 .
 (2) The H–O–H bond angle in H_2O is smaller than the H–N–H bond angle in NH_3 .
 (3) The H–C–H bond angle in CH_4 is larger than the H–N–H bond angle in NH_3 .
 (4) The H–C–H bond angle in CH_4 , the H–N–H bond angle in NH_3 , and the H–O–H bond angle in H_2O are all greater than 90°


Ans (1)

Molecules	Bond angle	Variations of bond angles
CH ₃	109° 28'	CH ₄ > NH ₃ > H ₂ O
$\ddot{\text{N}}\text{H}_3$	107°	
H ₂ O	104° 5'	$\text{H}-\text{C}-\text{H} > \text{H}-\text{N}-\text{H} > \text{H}-\text{O}-\text{H}$

20. Which one of the following statements is correct when SO₂ is passed through acidified K₂Cr₂O₇ solution

- (1) The solution is decolourized.
- (2) SO₂ is reduced.
- (3) Green Cr₂(SO₄)₃ is formed.
- (4) The solution turns blue.

Ans (3)



21. The correct thermodynamic conditions for the spontaneous reaction at all temperatures is

- | | |
|---------------------------------------|---------------------------------------|
| (1) $\Delta H > 0$ and $\Delta S < 0$ | (2) $\Delta H < 0$ and $\Delta S > 0$ |
| (3) $\Delta H < 0$ and $\Delta S < 0$ | (4) $\Delta H < 0$ and $\Delta S = 0$ |

Ans (2)

$\Delta H < 0$ [Exothermic]

$\Delta S > 0$ [Entropy increases]

$\Delta G < 0$

22. Natural rubber has

- (1) All trans-configuration
- (2) Alternate cis-and trans-configuration
- (3) Random cis-and trans-configuration
- (4) All cis-configuration

Ans (4)

Natural rubber is cis-polyisoprene

23. In which of the following options the order of arrangement does not agree with the variation of property indicated against it?

- (1) B < C < N < O (increasing first ionization enthalpy)
- (2) I < Br < Cl < F (increasing electron gain enthalpy)
- (3) Li < Na < K < Rb (increasing metallic radius)
- (4) Al³⁺ < Mg²⁺ < Na⁺ < F⁻ (increasing ionic size)

Ans (1)

	$\Delta H_E(\text{KJ/mol})$
O	1314
N	1402
C	1086
B	801

∴ The correct order for 1st ionization energy is B < C < O < N.

and

Ans (2)

	Electron gain enthalpy (KJ/mol)
F	-328
Cl	-349
Br	-325
I	-295

∴ The correct order for electron gain enthalpy is $I < Br < F < Cl$.

24. Which of the following reagents would distinguish cis-cyclopenta-1,2-diol from the trans-isomer?

- (1) Ozone (2) MnO_2
(3) Aluminium isopropoxide (4) Acetone


Ans (4)

Acetone gives ketal with cis isomer and not trans.

25. The product obtained as a result of a reaction of nitrogen with CaC_2 is

- (1) $CaCN$ (2) $CaCN_3$ (3) Ca_2CN (4) $Ca(CN)_2$

Ans (4)



26. Fog is a colloidal solution of

- (1) Gas in liquid (2) Solid in gas (3) Gas in gas (4) Liquid in gas

Ans (4)

fog is a colloid of water vapour in air.

27. Which one of the following orders is correct for the bond dissociation enthalpy of halogen molecules?

- (1) $Cl_2 > Br_2 > F_2 > I_2$ (2) $Br_2 > I_2 > F_2 > Cl_2$ (3) $F_2 > Cl_2 > Br_2 > I_2$ (4) $I_2 > Br_2 > Cl_2 > F_2$

Ans (1)

X_2	BDE (KJ/mol)
F_2	158
Cl_2	242
Br_2	193
I_2	151

The correct order of bond dissociation enthalpy is $Cl_2 > Br_2 > F_2 > I_2$.

28. Equal moles of hydrogen and oxygen gases are placed in a container with a pin-hole through which both can escape. What fraction of the oxygen escapes in the time required for one-half of the hydrogen to escape?

- (1) $\frac{1}{4}$ (2) $\frac{3}{8}$ (3) $\frac{1}{2}$ (4) $\frac{1}{8}$

Ans (4)

$$\frac{r_{H_2}}{r_{O_2}} = \sqrt{\frac{M_{O_2}}{M_{H_2}}} = \sqrt{\frac{32}{2}} = 4$$

$$r = \frac{V}{t} \quad \frac{1}{V_{O_2}} = 8$$

$$\frac{r_{H_2}}{r_{O_2}} = \frac{V_{H_2}/t}{V_{O_2}/t} \quad V_{O_2} = \frac{1}{8}$$

$$V_{H_2} = \frac{1}{2}$$

$$4 = \frac{1}{V_{O_2}}$$

29. Lithium has a bcc structure. Its density is 530 kg m^{-3} and its atomic mass is 6.94 g mol^{-1} . Calculate the edge length of a unit cell of Lithium metal. ($N_A = 6.02 \times 10^{23} \text{ mol}^{-1}$)

- (1) 352 pm (2) 527 pm (3) 264 pm (4) 154 pm

Ans (1)

$$d = \frac{Z.M}{a^3 N_A} = \frac{2 \times 6.94}{6.023 \times 10^{23} \times a^3}$$

$$a^3 = \frac{2 \times 6.94}{6.02 \times 10^{23} \times 0.530}$$

$$a = 352 \text{ pm}$$

30. Which of the following statements about the composition of the vapour over an ideal 1 : 1 molar mixture of benzene and toluene is correct? Assume that the temperature is constant at 25°C . (Given, Vapour Pressure Data at 25°C , benzene = 12.8 kPa, toluene = 3.85 kPa).

- (1) The vapour will contain a higher percentage of toluene
 (2) The vapour will contain equal amounts of benzene and toluene
 (3) Not enough information is given to make a prediction
 (4) The vapour will contain a higher percentage of benzene

Ans (4)

Benzene is more volatile.

31. Which of the following has longest C–O bond length? (Free C–O bond length in Co is 1.128 \AA)

- (1) $[\text{Co}(\text{CO})_4]^{2+}$ (2) $[\text{Fe}(\text{CO})_4]^{2-}$ (3) $[\text{Mn}(\text{CO})_6]^+$ (4) $\text{Ni}(\text{CO})_4$

Ans (3)


Octahedral complex have high π -back bonding hence C–O bond become longer.

32. Among the following, the correct order of acidity is

- (1) $\text{HClO} < \text{HClO}_2 < \text{HClO}_3 < \text{HClO}_4$ (2) $\text{HClO}_2 < \text{HClO} < \text{HClO}_3 < \text{HClO}_4$
 (3) $\text{HClO}_4 < \text{HClO}_2 < \text{HClO} < \text{HClO}_3$ (4) $\text{HClO}_3 < \text{HClO}_4 < \text{HClO}_2 < \text{HClO}$

Ans (1)


33. In the reaction



X and Y are

- (1) X = 2-Butyne ; Y = 3-Hexyne
 (2) X = 2-Butyne ; Y = 2-Hexyne
 (3) X = 1-Butyne ; Y = 2-Hexyne
 (4) X = 1-Butyne ; Y = 3-Hexyne

Ans (4)



34. MY and NY₃ two nearly insoluble salts, have the same K_{sp} values of 6.2 × 10⁻¹³ at room temperature. Which statement would be true in regard to MY and NY₃?
- (1) The molar solubility of MY in water is less than that of NY₃.
 - (2) The salts MY and NY₃ are more soluble in 0.5 M KY than in pure water.
 - (3) The addition of the salt of KY to solution of MY and NY₃ will have no effect on their solubilities.
 - (4) The molar solubilities of MY and NY₃ in water are identical.

Ans (1)

The solubility of MY is lower than NY₃

MY	NY ₃
$K_{sp} = s^2$	$K_{sp} = 27s^4$
$s = \sqrt{K_{sp}}$	$S = \sqrt[4]{\frac{K_{sp}}{27}}$


35. Consider the nitration of benzene using mixed conc. H₂SO₄ and HNO₃. If a large amount of KHSO₄ is added to the mixture, the rate of nitration will be
- (1) slower
 - (2) unchanged
 - (3) doubled
 - (4) faster

Ans (1)

Increase in concentration of HSO₄⁻, it retards the formation of the electrophile [NO₂⁺]

36. The product formed by the reaction of an aldehyde with a primary amine is
- (1) Ketone
 - (2) Carboxylic acid
 - (3) Aromatic acid
 - (4) Schiff base

Ans (4)



37. The pressure of H₂ required to make the potential of H₂-electrode zero in pure water at 298 K is
- (1) 10⁻¹² atm
 - (2) 10⁻¹⁰ atm
 - (3) 10⁻⁴ atm
 - (4) 10⁻¹⁴ atm

Ans (4)

38. The correct statement regarding RNA and DNA, respectively is
- (1) The sugar component in RNA is ribose and the sugar component in DNA is 2'-deoxyribose.
 - (2) The sugar component in RNA is arabinose and the sugar component in DNA is ribose.
 - (3) The sugar component in RNA is 2'-deoxyribose and the sugar component in DNA is arabinose.
 - (4) The sugar component in RNA is arabinose and the sugar component in DNA is 2'-deoxyribose.

Ans (1)

DNA – deoxy ribose sugar

RNA – Ribose sugar

39. Which one given below is a non-reducing sugar?
- (1) Lactose
 - (2) Glucose
 - (3) Sucrose
 - (4) Maltose

Ans (3)


Sucrose is the non reducing sugar due to C₁ – C₂ linkage between glucose and fructose respectively.

40. Which of the following statements about hydrogen is incorrect?
- (1) Hydrogen never acts as cation in ionic salts.
 - (2) Hydronium ion, H_3O^+ exists freely in solution.
 - (3) Dihydrogen does not act as a reducing agent.
 - (4) Hydrogen has three isotopes of which tritium is the most common.

Ans (4)

Most stable isotope of hydrogen is H_1^1 (protium)

41. Consider the following liquid – vapour equilibrium.







Which of the following relations is correct?

- (1) $\frac{d \ln P}{dT} = \frac{-\Delta H_v}{RT}$
- (2) $\frac{d \ln P}{dT^2} = \frac{-\Delta H_v}{T^2}$
- (3) $\frac{d \ln P}{dT} = \frac{\Delta H_v}{RT^2}$
- (4) $\frac{d \ln G}{dT^2} = \frac{\Delta H_v}{RT^2}$

Ans (3)

42. Which of the following biphenyls is optically active?



Ans (1)

43. Which of the following statements is false?

- (1) Ca^{2+} ions are important in blood clotting.
- (2) Ca^{2+} ions are not important in maintaining the regular beating of the heart.
- (3) Mg^{2+} ions are important in the green parts of plants.
- (4) Mg^{2+} ions form a complex with ATP.

Ans (2)

44. The ionic radii of A^+ and B^- ions are 0.98×10^{-10} m and 1.81×10^{-10} m. The coordination number of each ion in AB is:

- (1) 4
- (2) 8
- (3) 2
- (4) 6

Ans (4)

Radius **Ratio**

$$\frac{r_+}{r_-} = \frac{0.98 \times 10^{-10}}{1.81 \times 10^{-10}} = 0.5414$$

Hence cations occupy octahedral void with CN = 6

45. The rate of a first-order reaction is $0.04 \text{ mol l}^{-1} \text{ s}^{-1}$ at 10 seconds and $0.03 \text{ mol l}^{-1} \text{ s}^{-1}$ at 20 seconds after initiation of the reaction. The half-life period of the reaction is:

- (1) 34.1 s
- (2) 44.1 s
- (3) 54.1 s
- (4) 24.1 s

Ans (4)

$$K = \frac{2.303}{t} \log \left(\frac{C_1}{C_2} \right)$$

$$= \frac{2.303}{10} \log \left(\frac{3}{4} \right)$$

$$= 0.0278 \text{ min}^{-1}$$

$$t_{1/2} = \frac{0.693}{K}$$

$$= \frac{0.693}{0.0278} = 24.14 \text{ min}$$

46. The two polypeptides of human insulin are linked together by

- (1) phosphodiester bond
- (2) covalent bond
- (3) disulphide bridges
- (4) hydrogen bonds

Ans (3)

47. The coconut water from tender coconut represents

- (1) fleshy mesocarp
- (2) free nuclear proembryo
- (3) free nuclear endosperm
- (4) endocarp

Ans (3)

48. Which of the following is not a feature of the plasmids?

- (1) Circular structure
- (2) Transferable
- (3) Single-stranded
- (4) Independent replication

Ans (3)

Plasmids are always double stranded.

49. Which is the National Aquatic Animal of India?

- (1) River dolphin
- (2) Blue whale
- (3) Sea-horse
- (4) Gangetic shark

Ans (1)

Platanista gangetica [The Gangetic dolphin] was declared the National Aquatic Animal by Govt. of India.

50. The *Avena* curvature is used for bioassay of

- (1) GA₃
- (2) IAA
- (3) Ethylene
- (4) ABA

Ans (2)

The bioassay was done by Went.

51. Which of the following is the most important cause of animals and plants being driven to extinction?

- (1) Alien species invasion
- (2) Habitat loss and fragmentation
- (3) Co-extinctions
- (4) Over-exploitation

Ans (2)

Main cause

52. Which of the following approaches does not give the defined action of contraceptive?

(1)	Intra uterine devices	Increase phagocytosis of sperms, suppress sperm motility and fertilizing capacity of sperms
(2)	Hormonal contraceptives	Prevent/retard entry of sperms, prevent ovulation and fertilization
(3)	Vasectomy	Prevents spermatogenesis
(4)	Barrier methods	Prevent fertilization

Ans (3)

Vasectomy prevents sperms from entering ejaculatory duct.

53. In a testcross involving F_1 dihybrid flies, more parental-type offspring were produced than the recombinant-type offspring. This indicates

- (1) Chromosomes failed to separate during meiosis
- (2) The two genes are linked and present on the same chromosome
- (3) Both of the characters are controlled by more than one gene
- (4) The two genes are located on two different chromosomes

Ans (2)

The genes are linked

54. A typical fat molecule is made up of

- (1) One glycerol and three fatty acid molecules
- (2) One glycerol and one fatty acid molecule
- (3) Three glycerol and three fatty acid molecules
- (4) Three glycerol molecules and one fatty acid molecule

Ans (1)

55. Match the terms in Column I with description in Column II and choose the correct option.

Column I		Column II	
(a)	Dominance	(i)	Many genes govern a single character
(b)	Codominance	(ii)	In a heterozygous organism only one allele expresses itself
(c)	Pleiotropy	(iii)	In a heterozygous organism both alleles express themselves fully
(d)	Polygenic inheritance	(iv)	A single gene influences many characters

Code :

- | | | | |
|------------|------------|------------|------------|
| (a) | (b) | (c) | (d) |
| (1) (ii) | (iii) | (iv) | (i) |
| (2) (iv) | (i) | (ii) | (iii) |
| (3) (iv) | (iii) | (i) | (ii) |
| (4) (ii) | (i) | (iv) | (iii) |

Ans (1)

56. Which of the following statements is not correct?

- (1) Insects that consume pollen or nectar without bringing about pollination are called pollen/nectar robbers
- (2) Pollen germination and pollen tube growth are regulated by chemical components of pollen interacting with those the pistil
- (3) Some reptiles have also been reported as pollinators in some plant species
- (4) Pollen grains of many species can germinate on the stigma of a flower, but only one pollen tube of the same species grows into the style

Ans (4)

Because of pollen-pistil interaction.

57. Which of the following features is not present in *Periplaneta americana*?

- (1) Indeterminate and radial cleavage during embryonic development
- (2) Exoskeleton composed of N-acetylglucosamine
- (3) Metamerically segmented body
- (4) Schizocoelom as body cavity

Ans (1)

58. Water soluble pigments found in plant cell vacuoles are

- (1) Chlorophylls
- (2) Carotenoids
- (3) Anthocyanins
- (4) Xanthophylls

Ans (3)

Others are not water soluble.

59. A cell at telophase stage is observed by a student in a plant brought from the field. He tells his teacher that this cell is not like other cells at telophase stage. There is no formation of cell plate and thus the cell is containing more number of chromosomes as compared to other dividing cells. This would result in

- (1) Polyploidy
- (2) Somaclonal variation
- (3) Polyteny
- (4) Aneuploidy

Ans (1)

60. A plant in your garden avoids photorespiratory losses, has improved water use efficiency, shows high rates of photosynthesis at high temperatures and has improved efficiency of nitrogen utilisation. In which of the following physiological groups would you assign this plant?

- (1) C₄
- (2) CAM
- (3) Nitrogen fixer
- (4) C₃

Ans (1)

This is an adaptation for tropical high temperature regions.

61. In higher vertebrates, the immune system can distinguish self-cells and non-self. If this property is lost due to genetic abnormality and it attacks self-cells, then it leads to

- (1) Graft rejection
- (2) Auto-immune disease
- (3) Active immunity
- (4) Allergic response

Ans (2)

62. Emerson's enhancement effect and Red drop have been instrumental in the discovery of

- (1) Two photosystems operating simultaneously
- (2) Photophosphorylation and cyclic electron transport
- (3) Oxidative phosphorylation
- (4) Photophosphorylation and non-cyclic electron transport

Ans (1)

Discovery of PSI, PSII and their roles in photosynthesis.

63. Select the correct statement

- (1) *Salvinia*, *Ginkgo* and *Pinus* all are gymnosperms
- (2) *Sequoia* one of the tallest trees
- (3) The leaves of gymnosperms are not well adapted to extremes of climate
- (4) Gymnosperms are both homosporous and heterosporous

Ans (2)

64. Which of the following is not a characteristic feature during mitosis in somatic cells?

- (1) Disappearance of nucleolus
- (2) Chromosome movement
- (3) Synapsis
- (4) Spindle fibres

Ans (3)

65. Blood pressure in the pulmonary artery is

- (1) more than that in the carotid
- (2) more than that in the pulmonary vein
- (3) less than that in the venae cavae
- (4) same as that in the aorta

Ans (2)

66. Which of the following structures is homologous to the wing of a bird?

- (1) Wing of a Moth
- (2) Hind limb of Rabbit
- (3) Flipper of Whale
- (4) Dorsal fin of a Shark

Ans (3)

67. Seed formation without fertilization in flowering plants involves the process of

- (1) Budding
- (2) Somatic hybridization
- (3) Apomixis
- (4) Sporulation

Ans (3)

68. Name the chronic respiratory disorder caused mainly by cigarette smoking

- (1) Asthma
- (2) Respiratory acidosis
- (3) Respiratory alkalosis
- (4) Emphysema

Ans (4)

69. Spindle fibres attach on to

- (1) Kinetochore of the chromosome
- (2) Centromere of the chromosome
- (3) Kinetosome of the chromosome
- (4) Telomere of the chromosome

Ans (1)

70. In context of Amniocentesis, which of the following statement is incorrect?

- (1) It is used for prenatal sex determination
- (2) It can be used for detection of Down syndrome
- (3) It can be used for detection of Cleft palate
- (4) It is usually done when a woman is between 14-16 weeks pregnant

Ans (3)

71. Stems modified into flat green organs performing the functions of leaves are known as

- (1) Phyllodes (2) Phylloclades
(3) Scales (4) Cladodes

Ans (2)

72. In chloroplast the highest number of protons are found in

- (1) Lumen of thylakoids (2) Inter membrane space
(3) Antennae complex (4) Stroma

Ans (1)

Due to photolysis of H_2O , inflow of H^+ leading to chemiosmosis is ATP mann.

73. Nomenclature is governed by certain universal rules. Which one of the following is contrary to the rules of nomenclature?

- (1) The first word in a biological name represents the genus name, and the second is a specific epithet
(2) The names are written in Latin and are italicised
(3) When written by hand, the names are to be underlined
(4) Biological names can be written any language

Ans (4)

74. In meiosis crossing over is initiated at

- (1) Leptotene (2) Zygotene
(3) Diplotene (4) Pachytene

Ans (4)

75. Antivenom injection contains reformed antibodies while polio drops that are administered into the body contain

- (1) Harvested antibodies (2) Gamma globulin
(3) Attenuated pathogens (4) Activated pathogens

Ans (3)

In order to induce active immunity artificially.

76. The taq polymerase enzyme is obtained from

- (1) *Thiobacillus ferrooxidans* (2) *Bacillus subtilis*
(3) *Pseudomonas putida* (4) *Thermus aquaticus*

Ans (4)

77. Which of the following most appropriately describes haemophilia?

- (1) X-linked recessive gene disorder (2) Chromosomal disorder
(3) Dominant gene disorder (4) Recessive gene disorder

Ans (1)

78. The standard petal of a papilionaceous corolla is also called

- (1) Pappus (2) Vexillum (3) Corona (4) Carina

Ans (2)

Standard is also called vexillum.

79. Which part of the tobacco plant is infected by *Meloidogyne incognita*?
 (1) Leaf (2) Stem (3) Root (4) Flower

Ans (3)

80. Which of the following statements is wrong for viroids?
 (1) They are smaller than viruses (2) They cause infections
 (3) Their RNA is of high molecular weight (4) They lack a protein coat

Ans (3)

Viroids have low molecular weight RNA.

81. Which of the following statements is not true for cancer cells in relation to mutations?
 (1) Mutations destroy telomerase inhibitor
 (2) Mutations inactivate the cell control
 (3) Mutations inhibit production of telomerase
 (4) Mutations in proto-oncogenes accelerate the cell cycle

Ans (3)

82. Which type of tissue correctly matches with its location?

	Tissue	Location
(1)	Areolar tissue	Tendons
(2)	Transitional epithelium	Tip of nose
(3)	Cuboidal epithelium	Lining of stomach
(4)	Smooth muscle	Wall of intestine

Ans (4)

83. Which of the following pairs of hormones are not antagonistic (having effects) to each other?
 (1) Insulin – Glucagon
 (2) Aldosterone – Atrial Natriuretic Factor
 (3) Relaxin – Inhibin
 (4) Parathormone – Calcitonin

Ans (3)

84. Specialised epidermal cells surrounding the guard cells are called
 (1) Subsidiary cells (2) Bulliform cells
 (3) Lenticels (4) Complementary cells

Ans (1)

85. Fertilization in humans is practically feasible only if
 (1) the ovum and sperms are transported simultaneously to ampullary – isthmic junction of the fallopian tube
 (2) the ovum and sperms are transported simultaneously to ampullary – isthmic junction of the cervix
 (3) the sperms are transported into cervix within 48 hours of release of ovum in uterus
 (4) the sperms are transported into vagina just after the release of ovum in fallopian tube

Ans (1)

86. Which one of the following the starter codon?

- (1) UGA (2) UAA (3) UAG (4) AUG

Ans (4)

87. A river with an inflow of domestic sewage rich in organic waste may result in

- (1) increased population of aquatic food web organisms
(2) an increased production of fish due to biodegradable nutrients
(3) death of fish due to lack of oxygen
(4) drying of the river very soon due to algal bloom

Ans (3)

88. Following are the two statements regarding the origin of life

- (a) the earliest organisms that appeared on the earth were non-green and presumably anaerobes
(b) the first autotrophic organisms were the chemoautotrophs that never released oxygen.

Of the above statements which one of the following options is correct?

- (1) (b) is correct but (a) is false (2) both (a) and (b) are correct
(3) both (a) and (b) are false (4) (a) is correct but (b) is false

Ans (2)

89. A system of rotating crops with legume or grass pasture to improve soil structure and fertility is called

- (1) Contour farming
(2) Strip farming
(3) Shifting agriculture
(4) Ley farming

Ans (4)

To improve and balance soil fertility.

90. Gause's principle of competitive exclusion states that

- (1) Competition for the same resources excludes species having different food preferences
(2) No two species can occupy the same niche indefinitely for the same limiting resources
(3) Larger organism exclude smaller ones through competition
(4) More abundant species will exclude the less abundant species through competition

Ans (2)

91. Which of the following characteristic features always holds true for the corresponding group of animals?

(1)	Viviparous	Mammalia
(2)	Possess a mouth with an upper and a lower jaw	Chordata
(3)	3 - chambered heart with one incompletely divided ventricle	Reptilia
(4)	Cartilaginous endoskeleton	Chondrichthyes

Ans (4)

92. Changes in GnRH pulse frequency in females is controlled by circulating levels of

- (1) estrogen and inhibin (2) progesterone only
(3) progesterone and inhibin (4) estrogen and progesterone

Ans (4)

93. Microtubules are the constituents of
 (1) Spindle fibres, Centrioles and Cilia
 (2) Centrioles, Spindle fibres and Chromatin
 (3) Centrosome, Nucleosome and Centrioles
 (4) Cilia, Flagella and Peroxisomes
Ans (1)
94. Mitochondria and chloroplast are :
 (i) semi– autonomous organelles
 (ii) formed by division of pre-existing organelles and they contain DNA but lack protein synthesizing machinery.
 Which one of the following options is correct?
 (1) (i) is true but (ii) is false. (2) (i) is true but (ii) is false.
 (3) Both (i) and (ii) are false. (4) Both (i) and (ii) are correct
Ans (2)
 Presence of 70 s ribosomes helps in making some proteins
95. Photosensitive compound in human eye is made up of:
 (1) Opsin and Retinal (2) Opsin and Retinol
 (3) Transducin and Retinene (4) Guanosine and Retinol
Ans (1)
 Only after splitting, opsin and retinol are formed.
96. Chrysophytes, Euglenoids, Dinoflagellates and Slime moulds are included in the kingdom
 (1) Protista (2) Fungi (3) Animalia (4) Monera
Ans (1)
97. The primitive prokaryotes responsible for the production of biogas from the dung of ruminant animals, include the
 (1) Thermoacidophiles (2) Methanogens (3) Eubacteria (4) Halophiles
Ans (2)
98. Identify the correct statement on inhibin
 (1) Is produced by granulosa cells in ovary and inhibits the secretion of FSH.
 (2) Is produced by granulosa cells in ovary and inhibits the secretion of LH.
 (3) Is produced by nurse cells in testes and inhibits the secretion of LH.
 (4) Inhibits the secretion of LH, FSH and Prolactin.
Ans (1)
99. It is much easier for a small animal to run uphill than for a large animal, because
 (1) Smaller animals have a higher metabolic rate.
 (2) Small animals have a lower O₂ requirement.
 (3) The efficiency of muscles in large animals is less than in the small animals.
 (4) It is easier to carry a small body weight.
Ans (1)
 Surface area to volume ratio is high contributing to high metabolic rate

100. A tall true breeding garden pea plant is crossed with a dwarf true breeding garden pea plant. When the F_1 plants were selfed the resulting genotypes were in the ratio of
 (1) 1 : 2 : 1 :: Tall heterozygous : Tall homozygous : Dwarf
 (2) 3 : 1 :: Tall : Dwarf
 (3) 2 : 1 :: Dwarf : Tall
 (4) 1 : 2 : 1 :: Tall homozygous : Tall heterozygous : Dwarf
Ans (4)
101. Depletion of which gas in the atmosphere can lead to an increased incidence of skin cancers
 (1) Ozone (2) Ammonia (3) Methane (4) Nitrous oxide
Ans (1)
 UV rays increasingly reach earth
102. Which one of the following is a characteristic feature of cropland ecosystem?
 (1) Least genetic diversity (2) Absence of weeds
 (3) Ecological succession (4) Absence of soil organisms
Ans (1)
103. Tricarpellary, syncarpous gynoecium is found in flowers of
 (1) Solanaceae (2) Fabaceae (3) Poaceae (4) Liliaceae
Ans (4)
104. In which of the following, all three are macronutrients?
 (1) Iron, copper, molybdenum (2) Molybdenum, magnesium, manganese
 (3) Nitrogen, nickel, phosphorus (4) Boron, zinc, manganese
Ans
 No options are correctly matching
 All options include atleast 1 micronutrient.
105. Reduction in pH of blood will
 (1) reduce the blood supply to the brain.
 (2) decrease the affinity of hemoglobin with oxygen.
 (3) release bicarbonate ions by the liver.
 (4) reduce the rate of heart beat.
Ans (2)
 High H^+ concentration helps detach Hb from O_2 . This is called Bohr's effect
106. Lack of relaxation between successive stimuli in sustained muscle contraction is known as
 (1) Fatigue (2) Tetanus (3) Tonus (4) Spasm
Ans (2)
 Continued contraction is tetanus
107. Which one of the following statements is wrong?
 (1) Golden algae are also called desmids (2) Eubacteria are also called false bacteria.
 (3) Phycomycetes are also called algal fungi. (4) Cyanobacteria are also called blue-green algae.
Ans (2)

108. Which of the following is a restriction endonuclease?
 (1) Protease (2) DNase I (3) RNase (4) Hind II
Ans (4)
109. Which of the following would appear as the pioneer organisms on bare rocks?
 (1) Liverworts (2) Mosses (3) Green algae (4) Lichens
Ans (4)
110. Water vapour comes out from the plant leaf through the stomatal opening. Through the same stomatal opening carbon dioxide diffuses into the plant during photosynthesis. Reason out the above statements using one of following options
 (1) Both processes can happen together because the diffusion coefficient of water and CO₂ is different.
 (2) The above processes happen only during night time.
 (3) One process occurs during day time, and the other at night.
 (4) Both processes cannot happen simultaneously.
Ans (1)
111. Cotyledon of maize grain is called
 (1) coleorhiza (2) coleoptile (3) scutellum (4) plumule
Ans (3)
112. Which of the following guards the opening of hepatopancreatic duct into the duodenum?
 (1) Ileocaecal valve (2) Pyloric sphincter
 (3) Sphincter of Oddi (4) Semilunar valve
Ans (3)
113. In the stomach, gastric acid is secreted by the
 (1) parietal cells (2) peptic cells
 (3) acidic cells (4) gastrin secreting cells
Ans (1)
114. In mammals, which blood vessel would normally carry largest amount of urea?
 (1) Dorsal Aorta (2) Hepatic Vein
 (3) Hepatic Portal Vein (4) Renal Vein
Ans (2)
 As ornithine cycle in liver produces urea
115. The term ecosystem was coined by
 (1) A.G. Tansley (2) E. Haeckel
 (3) E. Warming (4) E.P. Odum
Ans (1)
116. Which of the following is required as inducer(s) for the expression of Lac operon?
 (1) galactose (2) lactose
 (3) lactose and galactose (4) glucose
Ans (2)

117. Which of the following is wrongly matched in the given table?

	Microbe	Product	Applicaiton
(1)	Monascus purpureus	Statins	Lowering of blood cholesterol
(2)	Streptococcus	Streptokinase	removal of clot from blood vessel
(3)	Clostridium butylicum	Lipase	Removal of oil stains
(4)	Trichoderma polysporum	Cyclosporin A	Immunosuppressive drug

Ans (3)

118. When does the growth rate of a population following the logistic model equal zero? The logistic models is given as $\frac{dN}{dt} = rN(1 - \frac{N}{K})$

(1) when N nears the carrying capacity of the habitat.

(1) when N / K equals zero

(3) when death rate is greater than birth rate.

(4) when N / K is exactly one.

Ans (4)

$$\frac{dN}{dt} = rN[1 - 1]$$

119. Which one of the following statements is not true?

(1) Exine of pollen grains is made up of sporopollenin

(2) Pollen grains of many species cause severe allergies

(3) Stored pollen in liquid nitrogen can be used in the crop breeding programmes

(4) Tapetum helps in the dehiscence of anther

Ans (4)

120. In bryophytes and pteridophytes, transport of male gametes requires

(1) Insects

(2) Birds

(3) Water

(4) Wind

Ans (3)

121. Which of the following is not a stem modification?

(1) Thorns of citrus

(2) Tendrils of cucumber

(3) Flattened structures of *Opuntia*

(4) Pitcher of *Nepenthes*

Ans (4)

122. Which one of the following cell organelles is enclosed by a single membrane?

(1) Chloroplasts

(2) Lysosomes

(3) Nuclei

(4) Mitochondria

Ans (2)

123. Analogous structures are a result of

(1) Convergent evolution

(2) Shared ancestry

(3) Stabilizing selection

(4) Divergent evolution

Ans (1)

124. Which one of the following statements is wrong?

(1) Cellulose is a polysaccharide.

(2) Uracil is a pyrimidine

(3) Glycine is a sulphur containing amino acid.

(4) Sucrose is a disaccharide.

Ans (3)

125. Proximal end of the filament of stamen is attached to the
 (1) Connective (2) Placenta (3) Thalamus or petal (4) Anther
Ans (3)
126. Which of the following is not required for any of the techniques of DNA fingerprinting available at present?
 (1) Zinc finger analysis (2) Restriction enzymes
 (3) DNA-DNA hybridization (4) Polymerase chain reaction
Ans (1)
127. Which one of the following characteristics is not shared by birds and mammals?
 (1) Breathing using lungs (2) Viviparity
 (3) Warm blooded nature (4) Ossified endoskeleton
Ans (2)
128. Select the incorrect statement
 (1) LH triggers ovulation in ovary.
 (2) LH and FSH decrease gradually during the follicular phase.
 (3) LH triggers secretion of androgens from the Leydig cells.
 (4) FSH stimulates the sertoli cells which help in spermiogenesis.
Ans (2)
129. The amino acid Tryptophan is the precursor for the synthesis of
 (1) Thyroxine and Triiodothyronine (2) Estrogen and Progesterone
 (3) Cortisol and Cortisone (4) Melatonin and Serotonin
Ans (4)
130. Joint Forest Management Concept was introduced in India during:
 (1) 1970 s (2) 1980 s (3) 1990 s (4) 1960 s
Ans (2)
131. One of the major components of cell wall of most fungus
 (1) Peptidoglycan (2) Cellulose (3) Hemicellulose (4) Chitin
Ans (4)
132. A complex of ribosomes attached to a single strand of RNA is known as
 (1) Polymer (2) Polypeptide (3) Okazaki fragment (4) Polysome
Ans (4)
133. Which of the following features is not present in the Phylum - Arthropoda?
 (1) Metameric segmentation (2) Parapodia
 (3) Jointed appendages (4) Chitinous exoskeleton
Ans (2)
134. Asthma may be attributed to
 (1) allergic reaction of the mast cells in the lungs
 (2) inflammation of the trachea
 (3) accumulation of fluid in the lungs
 (4) bacterial infection of the lungs
Ans (1)


135. Pick out the correct statements:

- (a) Haemophilia is a sex-linked recessive disease
 (b) Down's syndrome is due to aneuploidy.
 (c) Phenylketonuria is an autosomal recessive gene disorder.
 (d) Sickle cell anaemia is an X - linked recessive gene disorder.
- (1) (b) and (d) are correct. (2) (a), (c) and (d) are correct
 (3) (a), (b) and(c) are correct. (4) (a) and (d) are correct.

Ans (3)

136. A capacitor of $2 \mu\text{F}$ is charged as shown in the diagram. When the switch S is turned to position 2, the percentage of its stored energy dissipated is:

- (1) 20% (2) 75%
 (3) 80% (4) 0%




Ans (3)

$$U_i = \frac{1}{2} C_1 V^2 = \frac{1}{2} \times 2 \times 10^{-6} \times V^2 = 1 \times 10^{-6} V^2$$

$$\Delta U = \frac{1}{2} \frac{C_1 C_2}{C_1 + C_2} \cdot V^2 = \frac{1}{2} \times \frac{2 \times 8}{2 + 8} \times 10^{-6} V^2 = \frac{4}{5} \times 10^{-6} V^2 = 0.8 \times 10^{-6} V^2$$


loss of energy = 80%

137. To get output 1 for the following circuit, the correct choice for the input is:



- (1) $A = 1, B = 0, C = 0$ (2) $A = 1, B = 1, C = 0$
 (3) $A = 1, B = 0, C = 1$ (4) $A = 0, B = 1, C = 0$

Ans (3)



138. A potentiometer wire is 100 cm long and a constant potential difference is maintained across it, Two cells are connected in series first to support one another and then in opposite direction. The balance points are obtained at 50 cm and 10 cm from the positive end of the wire in the two cases. The ratio of emf 's is:

- (1) 5 : 4 (2) 3 : 4 (3) 3 : 2 (4) 5 : 1

Ans (3)

$$\frac{\epsilon_1 + \epsilon_2}{\epsilon_1 - \epsilon_2} = \frac{50}{10} = 5$$

$$\epsilon_1 + \epsilon_2 = 5\epsilon_1 - 5\epsilon_2$$

$$6\epsilon_2 = 4\epsilon_1$$

$$\frac{\epsilon_1}{\epsilon_2} = \frac{6}{4} = \frac{3}{2}$$

139. When a metallic surface is illuminated with radiation of wavelength λ , the stopping potential is V . If the same surface is illuminated with radiation of wavelength 2λ , the stopping potential is $\frac{V}{4}$. The threshold wavelength for the metallic surface is:

- (1) 5λ (2) $\frac{5}{2}\lambda$ (3) 3λ (4) 4λ

Ans (3)

$$\frac{hc}{\lambda} = \frac{hc}{\lambda_0} + eV$$

$$\frac{hc}{2\lambda} = \frac{hc}{\lambda_0} + e\left(\frac{V}{4}\right)$$

$$\frac{hc\left(\frac{1}{\lambda} - \frac{1}{\lambda_0}\right)}{hc\left(\frac{1}{2\lambda} - \frac{1}{\lambda_0}\right)} = \frac{eV}{\frac{eV}{4}}$$

$$\frac{\frac{\lambda_0 - \lambda}{\lambda\lambda_0}}{\frac{\lambda_0 - 2\lambda}{2\lambda\lambda_0}} = 4$$

$$\frac{\lambda_0 - \lambda}{\lambda\lambda_0} \cdot \frac{2\lambda\lambda_0}{\lambda_0 - 2\lambda} = 4$$

$$\frac{2(\lambda_0 - \lambda)}{\lambda_0 - 2\lambda} = 4$$

$$\lambda_0 - \lambda = 2\lambda_0 - 4\lambda$$

$$2\lambda_0 - \lambda_0 = 3\lambda$$

$$\lambda_0 = 3\lambda$$

140. Two non-mixing liquids of densities ρ and $n\rho$ ($n > 1$) are put in a container. The height of each liquid is h . A solid cylinder of length L and density d is put in this container. The cylinder floats with its axis vertical and length pL ($p < 1$) in the denser liquid. The density d is equal to:

- (1) $\{2 + (n+1)p\}\rho$ (2) $\{2 + (n-1)p\}\rho$ (3) $\{1 + (n-1)p\}\rho$ (4) $\{1 + (n+1)p\}\rho$

Ans (3)

$$V_b \rho_b g = V_b \rho_L g + V_b \rho'_L g$$


$$L\rho_b = (L - pL)\rho + pL(n\rho)$$

$$= L\rho - pL\rho + pLn\rho$$

$$\rho_b = \rho[L - pL + nPL]$$

$$\rho_b = \rho[1 - p + nP]$$

$$\rho_b = \rho[1 + P(n-1)]$$



141. Out of the following options which one can be used to produce a propagating electromagnetic wave?
- (1) A stationary charge (2) A charge less particle
 (3) An accelerating charge (4) A charge moving at constant velocity

Ans (3)

Accelerated charge produce EM waves

142. The charge flowing through a resistance R varies with time t as $Q = at - bt^2$, where a and b are positive constants. The total heat produced in R is:

- (1) $\frac{a^3R}{3b}$ (2) $\frac{a^3R}{2b}$ (3) $\frac{a^3R}{b}$ (4) $\frac{a^3R}{6b}$

Ans (4)

$$Q = at - bt^2$$

$$i = \frac{dq}{dt} = a - 2bt$$

$$Q \text{ for } t = \frac{a}{2b}, i = 0$$

$$\text{heat} = i^2Rdt$$

$$\int (a - 2bt) R dt$$

$$= \int a^2Rdt + 4b^2t^2Rdt - 4abt Rdt$$

$$= a^2R \cdot \int dt + 4b^2R \cdot \int t^2dt - 4abR \int t dt$$

$$a^2Rt \Big|_0^t + 4b^2R \cdot \frac{t^3}{3} \Big|_0^t - 4abR \cdot \frac{t^2}{2} \Big|_0^t$$

$$Q = a^2Rt + \frac{4b^2}{3}Rt^3 - \frac{4ab}{2}R \cdot t^2$$

$$\text{at } t = \frac{a}{2b}, i = 0$$

$$Q = \frac{a^2Ra}{2b} + \frac{4b^2}{3}R \frac{a^3}{8b^3} - \frac{2abR \cdot a^2}{4b^2}$$

$$= \frac{a^3R}{2b} + \frac{a^3R}{6b} - \frac{a^3R}{2b}$$

$$= \frac{a^3R}{6b}$$

143. At what height from the surface of earth the gravitation potential and the value of g are $-5.4 \times 10^7 \text{ J kg}^{-2}$ and 6.0 ms^{-2} respectively? Take the radius of earth as 6400 km:

- (1) 1600 km (2) 1400 km (3) 2000 km (4) 2600 km

Ans (4)

$$\frac{PE}{R+h} = g'$$

$$R+h = \frac{PE}{g'} = \frac{54 \times 10^6}{6}$$

$$h = 9 \times 10^6 - 6.4 \times 10^6$$

$$h = 2600 \text{ km}$$

144. Coefficient of linear expansion of brass and steel rods are α_1 and α_2 . Lengths of brass and steel rods are l_1 and l_2 respectively. If $(l_2 - l_1)$ is maintained same at all temperatures, which one of the following relations holds good?

- (1) $\alpha_1 l_2^2 = \alpha_2 l_1^2$ (2) $\alpha_1^2 l_2 = \alpha_2^2 l_1$ (3) $\alpha_1 l_1 = \alpha_2 l_2$ (4) $\alpha_1 l_2 = \alpha_2 l_1$

Ans (3)

$$\alpha_1 l_1 = \alpha_2 l_2$$

145. The intensity at the maximum in a Young's double slit experiment is I_0 . Distance between two slits is $d = 5\lambda$, where λ is the wavelength of light used in the experiment. What will be the intensity in front of one of the slits on the screen placed at a distance $D = 10d$?

- (1) $\frac{I_0}{4}$ (2) $\frac{3}{4}I_0$ (3) $\frac{I_0}{2}$ (4) I_0


Ans (3)

$$d \sin \theta = \frac{d5\lambda}{2 \times 50\lambda} = \frac{d}{20}$$

$$\Delta\phi = \frac{2\pi}{\lambda} \cdot \frac{5\lambda}{20} = \frac{\pi}{2}$$

$$\cos^2 \frac{\phi}{2} : \cos^2 \frac{\pi}{4}$$

$$I = \frac{I_0}{2}$$



146. Given the value of Rydberg constant is 10^7 m^{-1} , the wave number of the last line of the Balmer series in hydrogen spectrum will be:

- (1) $0.5 \times 10^7 \text{ m}^{-1}$ (2) $0.25 \times 10^7 \text{ m}^{-1}$ (3) $2.5 \times 10^7 \text{ m}^{-1}$ (4) $0.025 \times 10^4 \text{ m}^{-1}$

Ans (2)

$$\frac{1}{\lambda} = R \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right)$$

$$= R \left(\frac{1}{2^2} \right)$$

$$= \frac{R}{4}$$

$$= \frac{10^7}{4}$$

$$= 0.25 \times 10^7$$

147. The ratio of escape velocity at earth (v_e) to the escape velocity at a planet (v_p) whose radius and mean density are twice as that of earth is:

- (1) $1:2\sqrt{2}$ (2) $1:4$ (3) $1:\sqrt{2}$ (4) $1:2$

Ans (1)

$$v = \sqrt{\frac{2Gm}{R}}$$

$$= \sqrt{\frac{2G \cdot (dv)}{R}}$$

$$= \sqrt{\frac{2G}{R} \cdot dR^3}$$

$$v \propto \sqrt{R^2 \cdot d}$$

$$\propto R\sqrt{d}$$

$$\frac{v_1}{v_2} = \frac{R}{2R} \sqrt{\frac{d}{2d}} = \frac{1}{2\sqrt{2}}$$

$$\frac{v_1}{v_2} = \frac{1}{2\sqrt{2}}$$

148. A long solenoid has 1000 turns. When a current of 4A flows through it, the magnetic flux linked with each turn of the solenoid is 4×10^{-3} Wb. The self-inductance of the solenoid is:

- (1) 3 H (2) 2 H (3) 1 H (4) 4 H

Ans (3)

$$N = 1000; i = 4A, \phi = 4 \times 10^{-3}$$

$$L = \frac{N\phi}{i} = \frac{10^3 \times 4 \times 10^{-3}}{4} = 1H$$

149. A car is negotiating a curved road of radius R. The road is banked at an angle θ . The coefficient of friction between the tyres of the car and the road is μ_s . The maximum safe velocity on this road is:

- (1) $\sqrt{gR \frac{\mu_s + \tan \theta}{1 - \mu_s \tan \theta}}$ (2) $\sqrt{\frac{g}{R} \frac{\mu_s + \tan \theta}{1 - \mu_s \tan \theta}}$
 (3) $\sqrt{\frac{g}{R^2} \frac{\mu_s + \tan \theta}{1 - \mu_s \tan \theta}}$ (4) $\sqrt{gR^2 \frac{\mu_s + \tan \theta}{1 - \mu_s \tan \theta}}$

Ans (1)

$$V = \sqrt{gR \frac{\tan \theta + \mu_s}{1 - \mu_s \tan \theta}}$$

150. The magnetic susceptibility is negative for:

- (1) paramagnetic material only (2) ferromagnetic material only
 (3) paramagnetic and ferromagnetic material (4) diamagnetic material only

Ans (4)

χ is negative for diamagnetic substance.

151. A siren emitting a sound of frequency 800 Hz moves away from an observer towards a cliff at a speed of 15 ms^{-1} . Then, the frequency of sound that the observer hears in the echo reflected from the cliff is:

(Take velocity of sound in air = 330 ms^{-1})

- (1) 800 Hz (2) 838 Hz (3) 885 Hz (4) 765 Hz

Ans (2)

$$n' = \frac{v}{v-s} \cdot n$$

$$= \frac{330}{315}(800) = 838 \text{ Hz}$$

152. A body of mass 1 kg begins to move under the action of a time dependent force $\vec{F} = (2t\hat{i} + 3t^2\hat{j})\text{N}$, where \hat{i} and \hat{j} are unit vectors along x and y axis. What power will be developed by the force at the time t?

- (1) $(2t^2 + 4t^4) \text{ W}$ (2) $(2t^3 + 3t^4) \text{ W}$ (3) $(2t^3 + 3t^5) \text{ W}$ (4) $(2t^2 + 3t^3) \text{ W}$

Ans (3)

$$F = 2t\hat{i} + 3t^2\hat{j}$$

$$m \frac{dv}{dt} = 2t\hat{i} + 3t^2\hat{j}$$

$$\frac{dv}{dt} = 2t\hat{i} + 3t^2\hat{j}$$

$$dv = 2t\hat{i}dt + 3t^2\hat{j}dt$$

$$v = \frac{2t^2}{2}\hat{i} + 3\frac{t^3}{3}\hat{j} = t^2\hat{i} + t^3\hat{j}$$

$$p = F \cdot v = (2t^3 + 3t^5)W$$

153. From a disc of radius R and mass M , a circular hole of diameter R , whose rim passes through the centre is cut. What is the moment of inertia of the remaining part of the disc about a perpendicular axis, passing through the centre?
- (1) $13 MR^2 / 32$ (2) $11 MR^2 / 32$ (3) $9 MR^2 / 32$ (4) $15 MR^2 / 32$


Ans (1)

$$\frac{13MR^2}{32}$$

$$(MI_{YY})_{\text{Remaining}} = (MI_{yy})_{\text{whole}} - (MI_y)_{\text{hole}}$$

$$= \left(\frac{MR^2}{2}\right) - \left[\frac{M}{4} \cdot \left(\frac{R}{2}\right)^2\right] + \left(\frac{M}{4}\right)\left(\frac{R}{2}\right)^2$$

$$= \frac{MR^2}{2} - \left[\frac{MR^2}{32} + \frac{MR^2}{16}\right] = \frac{MR^2}{2} - \frac{3MR^2}{32} = \left(\frac{13MR^2}{32}\right)$$



154. In a diffraction pattern due to a single slit of width 'a', the first minimum is observed at an angle 30° when light of wavelength 5000 \AA is incident on the slit. The first secondary maximum is observed at an angle of:

- (1) $\sin^{-1}\left(\frac{2}{3}\right)$ (2) $\sin^{-1}\left(\frac{1}{2}\right)$ (3) $\sin^{-1}\left(\frac{3}{4}\right)$ (4) $\sin^{-1}\left(\frac{1}{4}\right)$

Ans (3)

$$a \sin \theta = \lambda$$

$$a \sin \theta = 5000 \text{ \AA}$$

$$\sin \theta = \frac{5000}{a} = \sin 30^\circ$$

$$\sin \theta_1 = \frac{3}{2}$$


$$\frac{\sin \theta_1}{\frac{1}{2}} = \frac{3 \frac{\lambda}{2a}}{\frac{5000}{a}}$$

$$\sin \theta_1 = \frac{3}{2} \times \frac{1}{2}$$

$$\theta_1 = \sin^{-1}\left(\frac{3}{4}\right)$$

155. A square loop ABCD carrying a current is placed near and coplanar with a long straight conductor XY carrying a current I , the net force on the loop will be:

- (1) $\frac{\mu_0 Ii}{2\pi}$ (2) $\frac{2\mu_0 IiL}{3\pi}$
- (3) $\frac{\mu_0 IiL}{2\pi}$ (4) $\frac{2\mu_0 Ii}{3\pi}$



Ans (4)

$$\frac{\mu_0 i I L}{2\pi L/2} - \frac{\mu_0 i I L}{2\pi \frac{3L}{2}}$$
$$\frac{\mu_0 i I L}{\pi L} \left[1 - \frac{1}{3} \right]$$
$$= \frac{2}{3} \frac{\mu_0 i I L}{\pi L} = \frac{2}{3} \frac{\mu_0 i I}{\pi}$$

156. A black body is at a temperature of 5760 K. The energy of radiation emitted by the body at wavelength 250 nm is U_1 , at wavelength 500 nm is U_2 and that at 1000 nm is U_3 . Wien's constant, $b = 2.88 \times 10^6$ nmK. Which of the following is correct?

- (1) $U_3 = 0$ (2) $U_1 > U_2$ (3) $U_2 > U_1$ (4) $U_1 = 0$

Ans (3)

$$\lambda_m T = b$$

$$\lambda_m(5760) = 2.88 \times 10^6$$

$$\lambda_m = \frac{2.88 \times 10^6}{5760} = 500 \text{ nm}$$

as λ_m is maximum = 500 nm

Corresponding energy is maximum $U_2 > U_1$

157. An air column, closed at one end and open at the other, resonate with a tuning fork when the smallest length of the column is 50 cm. The next larger length of the column resonating with the same tuning fork is:

- (1) 100 cm (2) 150 cm (3) 200 cm (4) 66.7 cm

Ans (2)

$$l_2 = 3l_1$$

resonating column

$$l_2 = 150 \text{ cm}$$

158. The molecules of a given mass of a gas have r.m.s. velocity of 200 ms^{-1} at 27°C and $1.0 \times 10^{-5} \text{ Nm}^{-2}$ pressure. When the temperature and pressure of the gas are respectively, 127°C and $0.05 \times 10^5 \text{ Nm}^{-2}$, the r.m.s. velocity of its molecules in ms^{-1} is:

- (1) $\frac{400}{\sqrt{3}}$ (2) $\frac{100\sqrt{2}}{3}$ (3) $\frac{100}{3}$ (4) $100\sqrt{2}$

Ans (1)

$$V_{\text{RMS}} = \sqrt{\frac{3RT}{n}}$$


$$V \propto \sqrt{T}$$

$$\frac{V_1}{V_2} = \sqrt{\frac{T_1}{T_2}} = \sqrt{\frac{300}{400}} = \frac{\sqrt{3}}{2}$$

$$2V_1 = \sqrt{3}V_2$$

$$V_2 = \frac{29}{\sqrt{3}} = \frac{2(200)}{\sqrt{3}} = \frac{400}{\sqrt{3}}$$

159. Consider the junction diode as ideal. The value of current flowing through AB is:



- (1) 10^{-2} A (2) 10^{-1} A (3) 10^{-3} A (4) 0 A

Ans (1)

$$i = \frac{10}{10^3} = 10^{-2} \text{ A}$$

160. If the magnitude of sum of two vectors is equal to the magnitude of difference of the two vectors, the angle between these vectors is:

- (1) 90° (2) 45° (3) 180° (4) 0°

Ans (1)

$$2F \cos\left(\frac{\theta}{2}\right) = 2F \sin\left(\frac{\theta}{2}\right)$$

$$\frac{\theta}{2} = 45^\circ$$

$$\theta = 90^\circ$$

161. A astronomical telescope has objective and eyepiece of focal lengths 40 cm and 4 cm respectively. To view an object 200 cm away from the objective, the lenses must be separated by a distance:

- (1) 46.0 cm (2) 50.0 cm (3) 54.0 cm (4) 37.3 cm

Ans (3)

$$f_0 = 40 \text{ cm}$$

$$U_0 = -200 \text{ cm}$$

$$\frac{1}{f_0} = \frac{1}{v_0} - \frac{1}{u_0}$$

$$\frac{1}{40} = \frac{1}{v_0} + \frac{1}{200} = \frac{5-1}{200} = \frac{4}{200}$$

$$v_0 = \frac{200}{4} = 50 \text{ cm}$$

$$L = v_0 + f_e$$

$$50 + 4 = 54 \text{ cm}$$

162. A npn transistor is connected in common emitter configuration in a given amplifier. A load resistance of 800Ω is connected in the collector circuit and the voltage drop across it is 0.8 V. If the current amplification factor is 0.96 and the input resistance of the circuit is 192Ω , the voltage gain and the power gain of the amplifier will respectively be:

- (1) 3.69, 3.84 (2) 4, 4 (3) 4, 3.69 (4) 4, 3.84

Ans (4)

$$\alpha = 0.96$$


$$\text{Voltage gain} = \alpha \frac{R_L}{R_i}$$

$$= (0.96) \frac{(800)}{192} = 4$$

$$\text{Power gain} = \alpha^2 \frac{R_L}{R_i} = 4(0.96) = 3.84$$

163. A gas is compressed isothermally to half its initial volume. The same gas is compressed separately through an adiabatic process until its volume is again reduced to half. Then:
- (1) Compressing the gas through adiabatic process will require more work to be done.
 - (2) Compressing the gas isothermally or adiabatically will require the same amount of work.
 - (3) Which of the case (whether compression through isothermal or through adiabatic process) requires more work will depend upon the atomicity of the gas.
 - (4) Compressing the gas isothermally will require more work to be done.

Ans (1)



Area under isothermal curve, so work done is more than adiabatic.

164. A long straight wire of radius a carries a steady current I . The current is uniformly distributed over its cross-section. The ratio of the magnetic fields B and B' , at radial distances $\frac{a}{2}$ and $2a$ respectively, from the axis of the wire is:

- (1) $\frac{1}{2}$ (2) 1 (3) 4 (4) $\frac{1}{4}$

Ans (2)

$$B_{in} = \frac{\mu_0 i r}{2\pi R^2}$$

$$B_2 = B_0 = \frac{\mu_0 i}{2\pi r}$$

Here $R = a$ $r = \frac{a}{2}$ $\therefore B_1 = \frac{\mu_0 i}{2\pi a^2} \cdot \frac{a}{2} = \frac{\mu_0 i}{4\pi a}$

Here $r = 2a$

$$B_2 = \frac{\mu_0 i}{2\pi \cdot 2a} = \frac{\mu_0 i}{4\pi a} \quad \therefore \frac{B_1}{B_2} = \frac{1}{1}$$

165. Match the corresponding entries of column 1 with column 2. [Where m is the magnification produced by the mirror]

	Column 1		Column 2
(A)	$m = -2$	(a)	Convex mirror
(B)	$m = -\frac{1}{2}$	(b)	Concave mirror
(C)	$m = +2$	(c)	Real image
(D)	$m = +\frac{1}{2}$	(d)	Virtual image

- (1) A \rightarrow a and c; B \rightarrow a and d; C \rightarrow a and b; D \rightarrow c and d
- (2) A \rightarrow a and d; B \rightarrow b and c; C \rightarrow b and d; D \rightarrow b and c
- (3) A \rightarrow c and d; B \rightarrow b and d; C \rightarrow b and c; D \rightarrow a and d
- (4) A \rightarrow b and c; B \rightarrow b and c; C \rightarrow b and d; D \rightarrow a and d

Ans (4)

Convex mirror m is +ve image is virtual

Concave mirror m is +ve virtual image m is -ve real image.

166. If the velocity of a particle is $v = At + Bt^2$, where A and B are constants, then the distance travelled by it between 1s and 2s is:

- (1) $3A + 7B$ (2) $\frac{3}{2}A + \frac{7}{3}B$ (3) $\frac{A}{2} + \frac{B}{3}$ (4) $\frac{3}{2}A + 4B$

Ans (2)

$$\frac{ds}{dt} = At + Bt^2$$

$$ds = At dt + Bt^2 dt$$

$$s = \frac{At^2}{2} + \frac{Bt^3}{3}$$

$$\frac{A}{2}(2^2 - 1^2) + \frac{B}{3}(2^3 - 1^3)$$

$$s = \frac{3A}{2} + \frac{7B}{3}$$

167. A disk and a sphere of same radius but different masses roll off on two inclined planes of the same altitude and length. Which one of the two objects gets to the bottom the plane first?

- (1) Sphere
(2) Both reach at the same time
(3) Depends on their masses
(4) Disk


Ans (1)

$$a = \frac{g \sin \theta}{1 + \frac{K^2}{R^2}}$$

$$s = \frac{1}{2}at^2$$

$$t = \sqrt{\frac{2s}{a}}$$

$$= \sqrt{\frac{2s}{\frac{g \sin \theta}{1 + K^2/R^2}}}$$



$$K \text{ for disk} = \frac{R}{\sqrt{2}}$$

$$K \text{ for sphere} = \sqrt{\frac{2}{5}}R$$

168. Two identical charged spheres suspended from a common point by two massless strings of lengths l , are initially at a distance d ($d \ll l$) apart because of their mutual repulsion. The charges begin to leak from both the spheres at a constant rate. As a result, the spheres approach each other with a velocity v . Then v varies as a function of the distance x between the spheres, as:

- (1) $v \propto x$ (2) $v \propto x^{-\frac{1}{2}}$ (3) $v \propto x^{-1}$ (4) $v \propto x^{\frac{1}{2}}$

Ans (2)

$$x^3 = \frac{q^2 l}{2\pi \epsilon_0 mg}$$

$$x^3 \propto q^2$$

$$3x^2 \frac{dx}{dt} \propto 2q \cdot \frac{dq}{dt}$$

$$3x^2 V \propto q$$

$$V \propto \frac{q}{x^2}$$

$$V \propto \frac{x^{\frac{3}{2}}}{x^2}$$

$$V \propto x^{-\frac{1}{2}}$$

169. A particle moves so that its position vector is given by $\vec{r} = \cos \omega t \hat{x} + \sin \omega t \hat{y}$. Where ω is a constant.

Which of the following is true?

- (1) Velocity and acceleration both are parallel to \vec{r} .
- (2) Velocity is perpendicular to \vec{r} and acceleration is directed towards to origin.
- (3) Velocity is perpendicular to \vec{r} and acceleration is directed away from the origin.
- (4) Velocity and acceleration both are perpendicular \vec{r} .

Ans (2)

$$\vec{r} = \cos \omega t \hat{i} + \sin \omega t \hat{j}$$

$$\vec{V} = \frac{d\vec{r}}{dt}$$

$$\vec{V} = [-\omega \sin \omega t \hat{i}] + \omega \cos \omega t \hat{j}$$

$$\vec{V} \cdot \vec{r} = 0$$

$\Rightarrow \vec{V}$ is perpendicular to \vec{r}

$$\frac{d\vec{V}}{dt} = \vec{a} = -\omega^2 \cos \omega t \hat{i} - \omega^2 \sin \omega t \hat{j}$$

$$\vec{a} = -\omega^2 [\vec{r}]$$

170. A piece of ice falls from a height h so that melts completely. Only one - quarter of the heat produced is absorbed by the ice and all energy of ice gets converted into heat during its fall. The value of h is:

[Latent heat of ice is 3.4×10^5 J/kg and $g = 10$ N/kg]

- (1) 544 km
- (2) 136 km
- (3) 68 km
- (4) 34 km

Ans (2)

$$\frac{1}{4} mgh = mL$$

$$h = \frac{4L}{g}$$

$$= \frac{4 \times 3.4 \times 10^5}{10}$$

$$= \frac{4 \times 34 \times 10^4}{10}$$

$$= 136 \text{ km}$$

171. A uniform circular disc of radius 50 cm at rest is free to turn about an axis which is perpendicular to its plane and passes through its centre. It is subjected to a torque which produces a constant angular acceleration of 2.0 rad s^{-2} . Its net acceleration in ms^{-2} at the end of 2.0 s is approximately:

- (1) 7.0
- (2) 6.0
- (3) 3.0
- (4) 8.0

Ans (4)

$$a_t = r\alpha = \frac{1}{2} \times 2 = 1 \text{ ms}^{-2}$$

$$\alpha t = \omega \Rightarrow \omega = 2 \times 2 = 4 \text{ rads}^{-1}$$

$$V = \omega r = 2 \text{ ms}^{-1}$$

$$a_c = \frac{v^2}{r} = \frac{4}{\left(\frac{1}{2}\right)} = 8 \text{ ms}^{-2} \text{ and } \sqrt{a_c^2 + a_r^2} = \sqrt{8^2 + 1^2} = \sqrt{65} \approx 8 \text{ ms}^{-2}$$

172. What is the minimum velocity with which a body of mass m must enter a vertical loop of radius R so that it can complete the loop?

- (1) $\sqrt{2gR}$ (2) $\sqrt{3gR}$ (3) $\sqrt{5gR}$ (4) \sqrt{gR}

Ans (3)

$$V_L = \sqrt{5gR}$$

173. A small signal voltage $V(t) = V_0 \sin \omega t$ is applied across an ideal capacitor C :

- (1) Over a full cycle the capacitor C does not consume any energy from the voltage source.
(2) Current $I(t)$ is in phase with voltage $V(t)$.
(3) Current $I(t)$ leads voltage $V(t)$ by 180° .
(4) Current $I(t)$, lags voltage $V(t)$ by 90° .

Ans (1)

Average power consumption in pure capacitor is zero.

174. A uniform rope of length L and mass m_1 hangs vertically from a rigid support. A block of mass m_2 is attached to the free end of the rope. A transverse pulse of wavelength λ_1 is produced at the lower end of the rope. The wavelength of the pulse when it reaches the top of the rope is λ_2 . The ratio $\frac{\lambda_2}{\lambda_1}$ is:

- (1) $\sqrt{\frac{m_1 + m_2}{m_2}}$ (2) $\sqrt{\frac{m_2}{m_1}}$ (3) $\sqrt{\frac{m_1 + m_2}{m_1}}$ (4) $\sqrt{\frac{m_1}{m_2}}$


Ans (1)

$$V = \sqrt{gx} = \lambda$$

$$x \propto \text{mass}$$

$$\sqrt{m_2} \propto \lambda_1; \lambda_2 \propto \sqrt{m_1 + m_2}$$

$$\frac{\lambda_2}{\lambda_1} = \sqrt{\frac{m_1 + m_2}{m_2}}$$



175. An inductor 20 mH , a capacitor $50 \text{ }\mu\text{F}$ and a resistor $40 \text{ }\Omega$ are connected in series across a source of emf $V = 10 \sin 340 t$. The power loss in A.C. circuit is:

- (1) 0.67 W (2) 0.76 W (3) 0.89 W (4) 0.51 W

Ans (4)

176. An electron of mass m and a photon have same energy E . The ratio of de-Broglie wavelengths associated with them is:

- (1) $\left(\frac{E}{2m}\right)^{\frac{1}{2}}$ (2) $c(2mE)^{\frac{1}{2}}$ (3) $\frac{1}{c}\left(\frac{2m}{E}\right)^{\frac{1}{2}}$ (4) $\frac{1}{c}\left(\frac{E}{2m}\right)^{\frac{1}{2}}$

(c being velocity of light)

Ans (4)

$$A_e = \frac{h}{p} = \frac{h}{\sqrt{2mE}}$$

$$E_p = \frac{hc}{\lambda_p}$$

$$X_p = \frac{hc}{E}$$

$$\begin{aligned} \Rightarrow \frac{\lambda_e}{\lambda_p} &= \frac{h}{\sqrt{2mE}} \cdot \frac{E}{hc} \\ &= \frac{1}{c} \sqrt{\frac{E}{2m}} \\ &= \frac{1}{c} \left(\frac{E}{2m} \right)^{\frac{1}{2}} \end{aligned}$$

177. When an α -particle of mass 'm' moving with velocity 'v' bombards on a heavy nucleus of charge 'Ze', its distance of closest approach from the nucleus depends on m as:

- (1) $\frac{1}{\sqrt{m}}$ (2) $\frac{1}{m^2}$ (3) m (4) $\frac{1}{m}$

Ans (4)

$$\frac{1}{2}mv^2 = \frac{1}{4\pi\epsilon_0} \cdot \frac{(2e)^2}{r} \quad r \propto \frac{1}{m}$$

178. A refrigerator works between 4°C and 30°C. It is required to remove 600 calories of heat every second in order to keep the temperature of the refrigerated space constant. The power required is:

(Take 1 cal = 4.2 Joules)

- (1) 23.65 W (2) 236.5 W (3) 2365 W (4) 2.365 W

Ans (2)

$$P = \frac{w}{t} = \frac{Q_1 - Q_2}{t} = \frac{Q_2 T_1}{T_2} - Q_2$$

$$Q_2 \left[\frac{T_1 - T_2}{T_2} \right]$$

$$= 600 \times 4.2 \left[\frac{303 - 277}{277} \right]$$

$$\frac{2520 \times 26}{277} = 236.5 \text{ w}$$

179. A particle of mass 10 g moves along a circle of radius 6.4 cm with a constant tangential acceleration. What is the magnitude of this acceleration if the kinetic energy of the particle becomes equal to 8×10^{-4} J by the end of the second revolution after the beginning of the motion?

- (1) 0.15 m/s² (2) 0.18 m/s² (3) 0.2 m/s² (4) 0.1 m/s²

Ans (4)

$$m = 10 \text{ gram}$$

$$r = 6.4 \text{ cm}$$

$$KE = 8 \times 10^{-4}$$

$$S = 4\pi r$$

$$ma \times 4\pi r = 8 \times 10^{-4}$$

$$10 \times 10^{-3} \times a \times 4\pi \times 6.4 = 8 \times 10^{-4} \times 10^{-2}$$

$$a = \frac{10^{-2} \times 8}{4\pi \times 6.4 \times 10^{-2}} \approx 0.1 \text{ ms}^{-1}$$

180. The angle of incidence for a ray of light at a refracting surface of a prism is 45° . The angle of prism is 60° . If the ray suffers minimum deviation through the prism, the angle of minimum deviation and refractive index of the material of the prism respectively, are:

(1) $30^\circ; \sqrt{2}$

(2) $45^\circ; \sqrt{2}$

(3) $30^\circ; \frac{1}{\sqrt{2}}$

(4) $45^\circ; \frac{1}{\sqrt{2}}$

Ans (1)

$$i_1 = 45^\circ$$

$$A = 60^\circ$$

$$A = 2r = 60$$

$$r = 30^\circ$$

$$n = \frac{\sin 45}{\sin 30} = \frac{\frac{1}{\sqrt{2}}}{\frac{1}{2}} = \sqrt{2}$$

$$d = 2i - A$$

$$= 2(45) - 60$$

$$= 30^\circ$$

* * *